

DISTRICT LEVEL HOUSEHOLD AND FACILITY SURVEY -4

DISTRICT FACT SHEET DHARWAD

(2012-13)



Introduction

Three rounds of District Level Household and Facility Surveys (DLHS) have been undertaken by the Ministry of Health and Family Welfare, Government of India in the past (Round-I in 1998-99, Round-II in 2002-04, and Round-III in 2007-08) with the main objective to provide reproductive and child health related database at district level in India. The data from these surveys have been useful in setting the benchmarks and examining the progress the country has made after the implementation of RCH programme. In addition, the evidences generated by these surveys have been useful for monitoring and evaluation of ongoing programmes and planning of suitable strategies by the central and state governments. In view of the completion of six years of National Rural Health Mission (2005-12), there was a felt need to focus on the achievements and improvements so far. The Ministry of Health and Family Welfare, Government of India, therefore initiated the process of conducting DLHS-4 during 2011-2012 and designated the International Institute for Population Sciences (IIPS) as the nodal agency to carry out the survey.

Bilingual questionnaires in the local language and English pertaining to Household, Clinical, Anthropometric and Bio-Chemical tests (CAB) and Ever Married Women (age 15-49) were used and canvassed using Computer Assisted Personal Interview (CAPI). The village and health facility questionnaires were canvassed by using paper & pen in DLHS-4. In the household questionnaire, information on all members of the household and socio-economic characteristics of the household, assets possessed, number of marriages, morbidities and deaths in the household since January 2008, etc. were collected. The ever-married women questionnaire contained information on women's characteristics, maternal care, immunization and childcare, contraception and fertility preferences, reproductive health including knowledge about HIV/AIDS. The village questionnaire contained information on availability of health, education and other facilities in the village, and whether the facilities are accessible throughout the year. The health facility questionnaire contained information on human resources, infrastructure, and services. For the first time, a population-linked facility survey has been conducted in DLHS-4. At the district level, all Community Health Centres and the District Hospitals and Sub Divisional Hospitals were covered. Further, all Sub-Health Centres and Primary Health Centres which were expected to serve the population of the selected PSUs were also covered. Fieldwork in Dharwad was conducted during December 2012 to June 2013, gathering information from 1,295 households, 1,190 ever married women and 48 health facilities.

This fact sheet gives information on the key indicators and trend for the district of Dharwad.

	DLHS-4 (2012-13)		DLHS-3 (2007-08)	
Indicators	TOTAL	RURAL	TOTAL	RURAL
Sample Size				
Households surveyed	1295	688	1257	568
Ever married women age15-49 years	1190	663	1246	596
Currently married women age 15-49 years interviewed	1086	611	1132	541
Population and household profile				
Percentage of population literate age 7+ years	79.3	74.2	75.3	69.7
Percentage of population below age 15 years	25.5	26.0	28.9	31.6
Mean household size	5.0	5.2	5.7	5.8
Sex ratio at birth (Male per 100 Female)	96.8	99.3	98.1	101.3
Percentage of households				
Having electricity	95.7	93.5	93.6	91.2
Improved source of drinking water ¹	91.7	87.7	86.0	74.2
Having access to improved toilet facility	55.5	33.8	47.9	21.3
Use clean fuel for cooking ²	38.8	9.1	25.8	2.8
Marriage				
Mean age at marriage for girls (marriages that occurred during the reference period)	20.2	19.5	19.6	19.4
Mean age at marriage for boys (marriages that occurred during the reference period)	26.3	25.7	27.2	27.2
Percentage of currently married women married below age 18 years (marriages that occurred during the reference period)	14.4	16.5	20.7	28.6
Percentage of currently married men married below age 21 years (marriages that occurred during the reference period)	4.9	5.5	5.0	6.5
Characteristics of women (%)				
Currently married women who are illiterate	31.3	37.7	35.5	49.3
Currently married women with 10 or more years of schooling	29.4	19.9	26.6	11.2
Fertility (%)				
Births to women aged 15-19 years out of total births ³	9.4	9.0	12.9	16.4
Women aged 20-24 years reporting birth order of 2 & above	55.3	59.0	46.2	56.8
Women aged 15-49 years who reported birth order of 3 & above	20.8	23.7	30.7	35.9
Women with two children wanting no more children	40.6	35.6	51.0	40.8
Mean no. of children ever born to women age 40-49 years	3.1	3.2	6.5	6.8
Current use of Family Planning Methods (%)				
Any method	62.2	64.8	62.6	63.8
Any modern method	62.5	65.3	61.7	63.5
Female sterilization	60.1	64.3	57.8	63.1
Male sterilization	0.5	0.2	0.3	0.2
Pill	0.7	0.5	0.9	0.2
IUD	0.4	0.3	1.9	0.0
Condom	.9	0.2	NA	NA
Unmet Need for Family Planning (%)				
Total unmet need ⁴	5.4	4.5	14.4	14.2
Unmet Need for spacing	2.8	2.1	8.9	9.5
Unmet Need for limiting	2.6	2.4	5.5	4.7

¹ Includes pipe water into dwelling, piped to yard/plot, public tap/standpipe/hand pump/tube well/bore well/well covered/protected spring, tanker/truck, cart with small tank/drum and packaged/bottled water. ² LPG/PNG/Electricity/Biogas, ³ DLHS-4 reference period is from 1-1-2008 to survey date; DLHS-3 reference period is from 1-1-2004 to survey date.

Unmet need for spacing includes fecund women who are neither pregnant nor amenorrhea, who are not using any method of family planning, and say they want to wait two or more years for their next birth. It also includes fecund women who are not using any method of family planning, and say they are unsure whether they want another child or who want another child but are unsure when to have the birth.

Unmet need for limiting includes fecund women who are neither pregnant nor amenorrhea, who are not using any method of family planning, and who want no more children (These definitions are similar to NFHS-3).

⁴Total unmet need refers to unmet need for limiting and spacing

	DLHS-4 (2012-13)		DLHS-3 (2007-08)	
Indicators		RURAL	TOTAL	RURAL
Quality of Family Planning Services (%)				
Non-users ever advised by health personnel to adopt any family planning method	45.3	42.5	17.1	18.3
Current users told about side effects of method		13.0	14.2	14.5
Users who received follow-up services for sterilization and IUD within 48 hours	76.2	76.2	96.0	96.4
Post-partum adoption of Family Planning for sterilization		96.5	93.1	93.6
Antenatal Care ⁵ (Women who had last live/still birth during reference period) (%)				
Pregnant women who received any antenatal check-up	97.2	98.3	91.6	93.1
Pregnant women who had antenatal check-up in first trimester		86.0	71.7	65.6
Pregnant women who had three or more ANC visits		90.6	81.0	78.6
Pregnant women who had at least one tetanus toxoid injection		99.4	90.3	91.1
Pregnant women whose Blood Pressure (BP) taken		93.2	91.0	92.3
Pregnant women who had blood tested (Hb)		92.1	9.1	92.5
Pregnant women whose abdomen examined		82.5	89.7	91.1
Pregnant women who consumed 100 or more IFA Tablets/Syrup equivalent		19.3	36.7	30.4
Pregnant women who had full antenatal care ⁵		19.3	27.7	23.4
Delivery Care (women who had live/still birth during reference period) (%)	10.2	10.0	21.1	20.4
Institutional delivery	93.5	92.8	66.6	55.0
Delivery at government health institutions		72.2	31.6	35.6
Delivery at private health institutions		20.6	34.9	19.4
Delivery by Caesarean section at government health institutions		7.8	4.2	6.9
Delivery by Caesarean section at private health institutions		7.8	7.8	3.1
Delivery at home				
		6.1	32.9	44.0
Delivery at home conducted by skilled health personnel (Out of total Deliveries)		1.1	9.1	10.1
Mothers who received post-natal care within 48 hours of Institutional delivery		87.8	77.9	72.5
Mothers who received post-natal care within two weeks of Institutional delivery		93.9	73.7	66.9
Delivery attended by skilled health personnel		93.9	75.7	64.7
Discharge of mothers from institution after minimum stay of 48 hours		80.2	69.6	61.9
Out of pocket expenditure per institutional delivery in Public health facility(Rs. in 000's)	2.27	2.12	NA	NA
Percentage of women who received JSY benefits	444	40.0		0.0
Home delivery		18.2	9.8	6.9
Institutional delivery	26.6	34.1	12.6	19.3
Percentage of Women who had				
Any Pregnancy complication ⁷		55.6	59.7	58.1
Any Delivery complication		42.2	47.0	44.7
Any Post-delivery complication ⁷		23.3	42.3	48.4
Problem of vaginal discharge during last three months		6.1	11.1	12.5
Menstrual related problems during last three months	7.5	6.5	15.6	16.9
Percentage of pregnancy resulted in				
Live Birth	94.5	93.9	95.1	94.4
Still Birth	1.0	.7	1.9	1.9
Induced abortion	1.3	1.4	1.2	1.4
Spontaneous abortion	3.1	4.0	1.9	2.3
Child Immunization (%) (Children age 12-23 months)				
Number of children	99	61	90	42
Received full vaccination ⁸	67.0	67.2	81.8	81.4
Received BCG vaccine	99.0	100.0	98.4	97.7
Received 3 doses of DPT vaccine	82.3	78.9	87.2	85.7
Received 3 doses of polio vaccine	87.5	89.7	92.1	90.7
Received measles vaccine	89.6	93.1	90.3	90.7
Children (age 9-35 months) received at least one dose of vitamin A supplement in last 6	82.3	83.5	67.8	61.8
months				

^{.5}Full ANC: At least three visits for antenatal check-up, one TT injection received and 100 IFA tablets or adequate amount of syrup consumed.

6Doctor/ANM/Nurse/midwife/LHV/Other health personnel, 7Women who had their last live/still birth since 1-1-2008. 8 BCG, 3-injection of DPT, 3 doses of Polio (excluding polio zero) and Measles

	DLHS-4 (2012-13)		DLHS-3 (2007-08)	
Indicators	TOTAL	RURAL	TOTAL	RURAL
Child feeding practices (based on last-born child in the reference period) (%)				
Children age 0-5 months exclusively breastfed ⁹	75.7	72.7	NA	NA
Children age 6-9 months receiving solid/semi-solid food and breast milk	57.9	45.5	78.8	83.3
Children age 12-23 months receiving breast feeding along with complementary feeding	63.2	64.9	NA	NA
Children age 6-35 months exclusively breastfed for at least 6 months	16.8	20.9	40.6	36.5
Children under 3 years breastfed within one hour of birth	62.3	60.6	40.8	36.3
Birth Weight (%) (age below 36 months)				
Percentage of Children weighed at birth	96.3	97.2	NA	NA
Percentage of Children with low birth weight (out of those who weighted) (below 2.5 kg)		19.7	NA	NA
Awareness about Diarrhoea (%)	10.0	10.7	1,7,1	1 47 4
Women know about what to do when a child gets diarrhoea	86.2	83.7	90.8	91.8
Awareness about ARI (%)	00.2	00	00.0	0.10
Women aware about danger signs of ARI ¹⁰	57.1	49.5	42.1	47.9
Treatment of childhood diseases (based on last two surviving children born during the	07.1	10.0	12.1	17.0
reference period) (%)				
Prevalence of diarrhoea in last 2 weeks for under 5 years old children	6.4	6.4	16.8	16.1
Children with diarrhoea in the last 2 weeks and received ORS ¹¹	60.7	62.5	59.0	53.6
Children with diarrhoea in the last 2 weeks and sought advice/treatment	92.6	86.7	72.7	59.3
Prevalence of ARI in last 2 weeks for under 5 years old children	7.4	6.8	8.1	8.2
Children with acute respiratory infection or fever in last 2 weeks and sought advice/treatment	96.8	100.0	86.3	83.3
Children with diarrhoea in the last 2 weeks given Zinc along with ORS	48.1	33.3	NA	NA
Awareness of RTI/STI and HIV/AIDS (%)				
Women who have heard of RTI/STI	51.1	45.4	43.6	39.4
Women who have heard of HIV/AIDS	78.2	73.8	90.3	88.4
Women who have any symptoms of RTI/STI	19.3	17.8	NA	NA
Women who know the place to go for testing of HIV/AIDS ¹²	88.0	87.4	77.0	74.2
Women underwent test for detecting HIV/AIDS ¹²	60.1	56.4	37.3	36.1
Utilization of Government Health Services (%)				
Antenatal care	69.9	76.1	33.1	54.9
Treatment for pregnancy complications	66.4	79.2	41.4	58.2
Treatment for post-delivery complications	55.6	73.3	40.2	43.1
Treatment for children with diarrhoea ¹³	53.8	57.1	NA	NA
Treatment for children with ARI ¹³	44.8	37.5	NA	NA
Birth Registration (%)				
Children below age 5 years having birth registration done	95.7	96.8	NA	NA
Children below age 5 years who received birth certificate (out of those registered)		93.0	NA	NA
Personal Habits (age 15 years and above) (%)				
Men who use any kind of smokeless tobacco	55.4	70.4	NA	NA
Women who use any kind of smokeless tobacco		41.0	NA	NA
Men who smoke		6.9	NA	NA
Women who smoke	0.3	0.2	NA	NA
Men who consume alcohol	8.5	7.9	NA	NA
Women who consume alcohol	0.2	0.1	NA	NA
⁹ Children Who were given nothing but breast milk till the survey date ¹⁰ Acute Respiratory Infections ¹¹ the women who have heard of HIV/AIDS. ¹³ Last two weeks	Oral Reh	ydration Solu		Based on
the women who have heard of HIV/AIDS. ¹³ Last two weeks				

	DLHS-4 (2012-13)		DLHS-3 (2007-08)	
Indicators	TOTAL	RURAL	TOTAL	RURAL
Reported Prevalence of Morbidity				
Any Injury	2.1	2.5	NA	NA
Acute Illness	6.6	5.6	NA	NA
Chronic Illness	5.0	3.4	NA	NA
Reported Prevalence of Chronic Illness during last one year (%)				
Disease of respiratory system	21.1	19.5	NA	NA
Disease of cardiovascular system	5.6	5.6	NA	NA
Persons suffering from tuberculosis	0.3	0.8	NA	NA
Anaemia Status by Haemoglobin Level ¹⁴ (%)				
Children (6-59 months) having anaemia	15.0	14.2	NA	NA
Children (6-59 months) having severe anaemia	15.0	14.2	NA	NA
Children (6-9 Years) having anaemia - Male	65.3	74.0	NA	NA
Children (6-9 Years) having severe anaemia - Male	2.5	2.0	NA	NA
Children (6-9 Years) having anaemia - Female	64.0	64.2	NA	NA
Children (6-9 Years) having severe anaemia - Female	0.9	1.5	NA	NA
Children (6-14 years) having anaemia - Male	57.6	58.5	NA	NA
Children (6-14 years) having severe anaemia - Male	3.5	3.0	NA	NA
Children (6-14 years) having anaemia - Female	63.2	61.8	NA	NA
Children (6-14 years) having severe anaemia - Female	3.8	2.8	NA	NA
Children (10-19 Years ¹⁵) having anaemia - Male	44.3	45.0	NA	NA
Children (10-19 Years ¹⁵) having severe anaemia - Male	3.9	2.6	NA	NA
Children (10-19 Years ¹⁵) having anaemia - Female	63.3	60.2	NA	NA
Children (10-19 Years ¹⁵) having severe anaemia - Female	5.1	5.0	NA	NA
Adolescents (15-19 years) having anaemia	52.3	52.4	NA	NA
Adolescents (15-19 years) having severe anaemia	4.6	5.2	NA	NA
Pregnant women (15-49 aged) having anaemia	66.0	73.6	NA	NA
Pregnant women (15-49 aged) having severe anaemia	3.8	5.7	NA	NA
Women (15-49 aged) having anaemia	68.0	72.5	NA	NA
Women (15-49 aged) having severe anaemia	5.8	5.4	NA	NA
Persons (20 years and above) having anaemia	52.3	56.3	NA	NA
Persons (20 years and above) having Severe anaemia	4.8	5.6	NA	NA
Blood Sugar Level (age 18 years and above) (%)				
Blood Sugar Level >140 mg/dl (high)	7.9	7.1	NA	NA
Blood Sugar Level >160 mg/dl (very high)	5.5	5.0	NA	NA
Hypertension (age 18 years and above) (%)				
Above Normal Range (Systolic >140 mm of Hg & Diastolic >90 mm of Hg)	19.5	15.6	NA	NA
Moderately High (Systolic >160 mm of Hg & Diastolic >100 mm of Hg)	7.4	4.9	NA	NA
Very High (Systolic >180 mm of Hg & Diastolic >110 mm of Hg)	2.3	1.3	NA	NA
lodized salt in Households (%)				
Households using iodized salt (15+ppm)	50.7	35.4	NA	NA
¹⁴ Any anaemia below 11g/dl, severe anaemia below 7g/dl. ¹⁵ Excluding age group 19 years Chronic Illness: Any person with symptoms persisting for longer than one month is defined as sufferir	ng from chro	onic illness		

	Number/Percentage	
Indicators	DLHS-4	DLHS-3
Villages covered		
Number of villages	25	22
Health Facilities covered		
Number of Sub-Health Centres	24	17
Number of Primary Health Centres (PHC)	20	11
Number of Community Health Centres (CHC) including Block PHC	NA	2
Number of Sub-Divisional Hospitals (SDH)	3	NA
Number of District Hospitals (DH)	1	1
Health programmes at village level		
Percentage of villages having ASHA	96.0	9.1
Percentage of Villages having Village Health Nutrition and Sanitation Committee (VHNSC)	76.0	63.6
Accessibility of health facility (%)		
Villages with Sub-Health Centre within 3 km	80.0	68.2
Villages with PHC within 10 km	76.0	63.6
Availability of Health Infrastructure, Staff and Services at (%)		
Sub-Health Centre		
Sub-Health Centre located in government building	54.2	58.8
Sub-Health Centre located in government building	100.0	94.1
Sub-Health Centre with male health worker	62.5	70.6
Sub-Health Centre with ANM residing in Sub-Health Centre quarter where facility is available	58.3	77.8
Sub-Health Centre with additional ANM	0.0	0.0
Primary Health Centre (PHC)	0.0	0.0
PHCs functioning on 24 X 7 hours basis	75.0	54.6
PHCs having Lady Medical Officer	5.0	18.2
PHCs with at least 4 beds	100.0	81.8
PHCs with AYUSH doctor	60.0	27.3
PHCs having residential quarter for Medical Officer	55.0	54.6
PHCs having new born care services on 24 X 7 hours basis	100.0	NA
PHCs having referral services for pregnancies/delivery on 24 X 7 hours basis	93.3	NA
PHCs conducted at least 10 deliveries during last one month on 24 X 7 hours basis	86.7	18.2
Community Health Centre (CHC)		
CHCs having 24 X 7 hours normal delivery services	NA	2
CHCs having Obstetrician/Gynaecologist	NA	0
CHCs having Anesthetist	NA	Na
CHCs having functional Operation Theatre	NA	2
CHCs designated as FRUs	NA	2
CHCs designated as FRUs offering caesarean section	NA	0
CHCs having new born care services on 24 X 7 hours basis	NA	1
Sub Divisional Hospital (SDH)		
SDHs having Paediatrician	1	NA
SDHs having regular radiographer	0	NA
SDHs having 2D Echo facility	0	NA
SDHs having Ultrasound facility	0	NA
SDHs having three phase connection	3	NA
SDHs having critical care area	1	NA
SDHs having suggestion and complaint box	2	NA
District Hospital (DH)		
DHs having Paediatrician	1	1
DHs having regular radiographer	0	0
DHs having 2D Echo facility	0	0
DHs having Ultrasound facility	1	Na
DHs having three phase connection	1	1
DHs having critical care area	1	0
DHs having suggestion and complaint box	1	1
U	•	•

INTERNATIONAL INSTITUTE FOR POPULATION SCIENCES

Vision: "To position IIPS as a premier teaching and research institution in population sciences responsive to

emerging national and global needs based on values of inclusion, sensitivity and rights protection."

Mission: "The Institute will strive to be a centre of excellence on population, health and development issues

through high quality education, teaching and research. This will be achieved by (a) creating competent professionals, (b) generating and disseminating scientific knowledge and evidence, (c) collaboration

and exchange of knowledge, and (d) advocacy and awareness."

For additional information, please contact:

Director/Project Coordinator (DLHS-4) International Institute for Population Sciences

Govandi Station Road, Deonar Mumbai - 400 088 (India)

Telephone: 022-2556 3254/5/6, 022-4237 2465, 42372411

Fax: 022-25563257, 25555895

Email: rchpro@iips.net, director@iips.net

Website: http://www.rchiips.org

http://www.iipsindia.org

Additional Director General (Stat.) Ministry of Health and Family Welfare

Government of India Nirman Bhavan New Delhi 110 011

Telephone: 011 - 23061334

Fax: 011 - 23061334

Email: adg-mohfw@nic.in

Chief Director (Stat.)

Ministry of Health and Family Welfare

Government of India

Nirman Bhavan

New Delhi 110 011

Telephone: 011 - 23062699

Fax: 011 - 23062699 Email: cdstat@nic.in

Website: http://www.mohfw.nic.in